

## Reference

E-40

1. NCD for Neuromuscular Electrical Stimulation (NMES)(160.12).
2. Carty A, McCormack K, Coughlan GF, et al. Alterations in body composition and spasticity following subtetanic neuromuscular electrical stimulation training in spinal cord injury. *Journal of Rehabilitation Research & Development*. 2013;50:193-202.
3. Bethoux F, Rogers L, Nolan K, et al. Long-term follow-up to a randomized controlled trial comparing peroneal nerve functional electrical stimulation to an ankle foot orthosis for patients with chronic stroke. *Neurorehabil Neural Repair*. 2015;29(10):911-922
4. Mayer L, Warring T, Agrella S, et al. Effects of functional electrical stimulation on gait function and quality of life for people with multiple sclerosis taking dalfampridine. *Int J MS Care*. 2015;17:35–41.
5. Hayes, Inc. Hayes Medical Technology Directory Report. *Functional Electrical Stimulation (FES) for Treatment of Foot Drop in Multiple Sclerosis Patients*. Lansdale, PA: Hayes, Inc. July 16, 2015. Accessed May 11, 2018.
6. Hayes, Inc. Hayes Medical Technology Directory Report. *Functional Electrical Stimulation (FES) for Upper Extremity Rehabilitation Post Stroke*. Lansdale, PA: Hayes, Inc. July 8, 2009. Accessed May 11, 2018.
7. Hayes Health Technology Brief. *Functional Electrical stimulation for Rehabilitation Following Spinal Cord Injury*. Lansdale, PA: Hayes, Inc. November 16, 2017. Accessed May 11, 2018.
8. Michael G. Fehlings, MD, PhD, FRCSC, FACS et al. A Clinical Practice Guideline for the Management of Patients with Acute Spinal Cord Injury: Recommendations on the Type and Timing of Rehabilitation. *Global Spine Journal*. 2017;7(2)231S-238S.