

References

E-7

1. Clinical Trial- Centre for Research and Implementation of Clinical Practice in the U.K. and Tactile Technologies.
2. Dolibog P, Franek A, Taradaj J, et al. A comparative clinical study on five types of compression therapy in patients with venous leg ulcers. *Int J Med Sci.* 2013;11(1):34-43.
3. International Society of Lymphology. The diagnosis and treatment of peripheral lymphedema: 2013 Consensus Document of the International Society of Lymphology. *Lymphology.* 2013; 46(1):1-11.
4. Muluk SC, Hirsch AT, Taffe EC. Pneumatic compression device treatment of lower extremity lymphedema elicits improved limb volume and patient-reported outcomes. *Eur J Vasc Endovasc Surg.* 2013;46(4):480-487.
5. Mohler ER, Mondry TE. Prevention and treatment of lymphedema. *UpToDate.* Literature review current through: May 2014.
6. American College of Obstetricians and Gynecologists. Prevention of deep vein thrombosis and pulmonary embolism. Washington (DC). American College of Obstetricians and Gynecologists (ACOG); 2007 Aug. Reaffirmed in 2012.
7. Jacobs JJ, Watters III WC, Mont MA, et al. Preventing venous thromboembolic disease in patients undergoing elective hip and knee arthroplasty. American Academy of Orthopedic Surgeons (AAOS). Rosemont, IL. 2011.
8. Falck-Ytter Y, Francis CW, Johanson NA, et al. Prevention of VTE in orthopedic surgery patients: Antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest.* 2012;141(2 Suppl): e278S-325S.
9. Colwell CW Jr, Froimson MI, Anseth SD, et al. A mobile compression device for thrombosis prevention in hip and knee arthroplasty. *J Bone Joint Surg Am.* 2014;96(3):177-183.
10. Koo KH, Choi JS, Ahn JH, et al. Comparison of clinical and physiological efficacies of different intermittent sequential pneumatic compression devices in preventing deep vein thrombosis: A prospective randomized study. *Clinics in Orthopedic Surgery.* 2014;6:468-475.
11. Karaca-Mandic P, Hirsch AT, Rockson SG, et al. The cutaneous, net clinical, and health economic benefits of advanced pneumatic compression devices in patients with lymphedema. *JAMA Dermatol.* 2015;151(11):1187-1193.
12. CLOTS (Clots in Legs or stockings after Stroke) Trials Collaboration. "Effectiveness of intermittent pneumatic compression in reduction of risk of deep vein thrombosis in patients who have had a stroke (CLOTS 3): a multicentre randomised controlled trial." *The Lancet.* 2013;382(9891); 516-524.
13. Moran P, Teljeur C, Harrington P, Ryan M. A systematic review of intermittent pneumatic compression for critical limb ischaemia. *Vascular Medicine (London, England) [serial online].* 2015;20(1):41-50.
14. Hayes, Inc, Hayes Health Technology Brief. *Flexitouch System (tactile Systems Technology Inc.) for Lymphedema.* Lansdale, PA: Hayes, Inc.: October, 2107.

15. Hayes, Inc, Hayes Medical Technology Directory Report. *Pneumatic Compression for Prevention of Deep Vein Thrombosis Following Hip Surgery* . Lansdale, PA: Hayes, Inc.: Published May 2011, reviewed April 2015.
16. Hayes, Inc, Hayes Medical Technology Directory Report. *Intermittent Pneumatic Compression for Peripheral Arterial Disease*. Lansdale, PA: Hayes, Inc.: July 2013, reviewed June 9, 2017.