

## References

### L-102

1. National Opioid Use Guideline Group (NOUGG): Canada. Canadian Guideline for Safe and Effective Use of Opioids for Chronic Non-Cancer Pain. 2010.
2. Owen G, Burton A, Schade C, et al. Urine drug testing: current recommendations and best practices. *Pain Physician*. 2012; 15(3):ES119-133.
3. Manchikanti L, Abdi S, Atluri S, et al. American Society of Interventional Pain Physicians (ASIPP) guidelines for responsible opioid prescribing in chronic non-cancer pain: Part 2-guidance. *Pain Physician*. 2012; 15(3 Suppl.):S67-116.
4. Hamill-Ruth R, Larriviere K, McMasters M. Addition of objective data to identify risk for medication misuse and abuse: the inconsistency score. *Pain Medicine*. 2013; 14(12):1900-1907.
5. Gudín J, Moqali S, Jones J, et al. Risks, management, and monitoring of combination opioid, benzodiazepines, and/or alcohol use. *Postgrad Medicine*. 2013; 125(4):115-130.
6. Melanson S, Ptolemy A, Wasan A. Optimizing urine drug testing for monitoring medications compliance in pain management. *Pain Medicine*. 2013; 14(12):1813-1820.
7. Washington State Agency Medical Directors' Group. Interagency guideline on opioid dosing for chronic non-cancer pain: An educational aid to improve care and safety with opioid treatment. 2010.
8. Ward M, Hackenmueller S, Strathmann F, et al. Pathology consultation on urine compliance testing and drug abuse screening. *American Journal of Clinical Pathology*. 2014; 142(5):586-93.
9. Nuckols T, Anderson L, Popescu I, et al. Opioid prescribing: A systematic review and critical appraisal of guidelines for chronic pain. *Annals of Internal Medicine*. 2014; 160(1):38-47.
10. American Society of Addiction Medicine (ASAM). Public Policy Statement on Drug Testing as a Component of Addiction Treatment and Monitoring Programs and in other Clinical Settings. 2010.
11. Dupouy J, Memier V, Catala H, et al. Does urine drug abuse screening help for managing patients? A systematic review. *Drug Alcohol Depend*. 2014; 136:11-20.
12. Conermann T, Gosalia A, Kabazie A, et al. Utility of oral fluid in compliance monitoring of opioid medications. *Pain Physician*. 2014; 17(1):63-70.
13. American College of Occupational and Environmental Medicine (ACOEM). Principles for ensuring the safe management of pain medication prescriptions by occupational and environmental medicine physicians. 2012.
14. American Pain Society (APS)/ American Academy of Pain Medicine (AAPM). Clinical guidelines for chronic opioid therapy in patients with noncancer pain. 2013.
15. Davis K, Sadler A, Genzen J. A retrospective analysis of urine drugs of abuse immunoassay true positive rates at a national reference laboratory. *Journal of Analytical Toxicology*. 2016; 40(2):97-107.
16. Krishnamurthy P, Ranganathan G, Williams C, et al. Impact of urine drug screening on no shows and dropouts among chronic pain patients: A propensity-matched cohort study. *Pain Physician*. 2016; 19(2):89-100.

17. Dowell D, Haegerich TM, Chou R. Centers for Disease Control and Prevention (CDC) guideline for prescribing opioids for chronic pain-United States, 2016. *Journal of the American Medical Association*. 2016; 315(15):1624-1645.
18. Washington State Agency Medical Directors' Group. Interagency guideline on prescribing opioid dosing for pain. 2015.