

References

L-114

1. Sachse C, Brockmüller J, Bauer S, Roots I. Cytochrome P450 2D6 variants in a Caucasian population: allele frequencies and phenotypic consequences. *Am J Hum Genetics*. 1997;60:284-95.
2. Schroth W, Antoniadou L, Fritz P, et al. Breast cancer treatment outcome with adjuvant tamoxifen relative to patient CYP2D6 and CYP2C19 genotypes. *J Clin Oncol*. 2007;25(33):5187-93.
3. Goetz MP, Knox SK, Suman VJ, et al. The impact of cytochrome P450 2D6 metabolism in women receiving adjuvant tamoxifen. *Breast Cancer Res Treat*. 2007;101:113-21.
4. Newman WG, Hadfield KD, Latif A, et al. Impaired tamoxifen metabolism reduces survival in familial breast cancer patients. *Clin Cancer Res*. 2008;14(18):5913-8.
5. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Breast Cancer, Version 2.2016.
6. Visvanathan K, Lippman SM, Hurley P, Temin S. American Society of Clinical Oncology clinical practice guideline update on the use of pharmacologic interventions including tamoxifen, raloxifene, and aromatase inhibition for breast cancer risk reduction. *Gynecol Oncol*. 2009;115(1):132-4.
7. Schroth W, et al. Association between CYP2D6 polymorphisms and outcomes among women with early stage breast cancer treated with tamoxifen. *JAMA*. 2009;302(13):1429-3.
8. Seruga B, Amir E. Cytochrome P450 2D6 and outcomes of adjuvant tamoxifen therapy: results of a meta-analysis. *Breast Cancer Res Treat*. 2010 Aug; 122(3):609-17.
9. Companion diagnostics. U.S. Food & Drug Administration website.
10. Scheerens H, Malong A, Bassett K, et al Current status of companion and complementary diagnostics. *Clin Transl Sci*. 2017;10:84-92.