

## References

M-76

1. Mercuri A, Sfara C, Cavazzoni E, et al. Anal Cancer Screening for Diagnosis and Treatment of Anal Squamous Intraepithelial Lesions in HIV-positive MSM Patients. *J AIDS Clin Res* 2013, 5:1.
2. Hillman RJ, Cuming T, Darragh T, et al. 2016 IANS International Guidelines for Practice Standards in the Detection of Anal Cancer Precursors. *Journal of Lower Genital Tract Disease*. 2016;283-291.
3. Goon P, Morrison V, Fearnhead N, et al. High resolution anoscopy may be useful in achieving reductions in anal cancer local disease failure rates. *European Journal of Cancer Care*. 2015;411-416.
4. Hillman RJ, Gunathilake MPW, Jin F, et al. Ability to detect high-grade squamous anal intraepithelial lesions at high resolution anoscopy improves over time. *Sexual Health*. 2016;177–181.
5. Hayes, Inc. Hayes Medical Technology Directory Report. *High-Resolution Anoscopy for the Evaluation of Anal Lesions*. Lansdale,PA: Hayes, Inc; Published October 11, 2017. Reviewed November 6, 2014. Accessed on June 22, 2018.
6. Dalla Pria A, Alfa-Wali M, Fox P, et al. High-resolution anoscopy screening of HIV-positive MSM: Longitudinal results from a pilot study. *AIDS*. 2014;28(6):861-867.
7. Schofield AM, Sadler L, Nelson L, et al. A prospective study of anal cancer screening in HIV-positive and negative MSM. *AIDS*. 2016;30(9):1375-1383.