

References

S-147

1. Ibarra-Drendall C, Wilke L, Zalles C, et al. Reproducibility of Random periareolar fine needle aspiration in a multi-institutional cancer and leukemia group B (CALGB) cross-sectional study. *Cancer-Epidemiol Biomarkers*. 2009;18(5):1379-1385.
2. Dubowy A, Raubach M, Topalidis T, et al. Breast Duct Endoscopy: Ductoscopy from a diagnostic to an interventional procedure and its future perspective. *Acta Chir Belg*. 2011;(111):142-145.
3. Konstandiadou I, Mastoraki A, Kotsilianou, et al. Does ductal lavage assert its role as a noninvasive diagnostic modality to identify women at low risk of breast cancer development. *J Gynecol Oncol*. 23(2):110-114.
4. Tang S, Twelves D, Isacke C, Gui G. Mammary ductoscopy in the current management of breast disease. *Surg Endosc*. 2011; 25:1712-1722.
5. Hoffman A, Pellenberg R, Drendall C, et al. Comparison of Random periareolar fine needle aspirate versus ductal lavage for risk assessment and prevention of breast cancer. *Curr Breast Cancer Rep*. 2012;(4)(3):180-187.
6. Han Y, Li J, Han S, Jia S, Zhang Y, Zhang W. Diagnostic value of endoscopic appearance during ductoscopy in patients with pathological nipple discharge. *BMC Cancer*. May 2, 2017;17(1):300.
7. Do Canto L, Marian C, Haddad B, et al. MicroRNA analysis of breast ductal fluid in breast cancer patients. *International Journal Of Oncology*. May 2016;48(5):2071-2078.
8. Matos Do Canto L, Marian C, Haddad B, et al. Metabolomic profiling of breast tumors using ductal fluid. *International Journal Of Oncology*. December 2016;49(6):2245-2254.