

References

S-176

1. Jackson TJ, Domb BG, Stake CE, et al. Revision Hip Preservation Surgery with Hip Arthroscopy: Clinical outcomes. Arthroscopy Association of North America. 2014 Mar 15.
2. Tranovich MJ, Salzler MJ, Enseki KR, et al. A review of femoroacetabular impingement and hip arthroscopy in the athlete. *The Physician and Sports Medicine*. 2014;42(1):75-87.
3. Gupta AK, Abrams GD, Nho SJ. What's new in femoroacetabular impingement surgery: Will we be better in 2023? *Sports Health*. 2014;6(2):162-170.
4. Loudon JK, Reiman MP. Conservative management of femoroacetabular impingement (FAI) in the long distance runner. *Physical Therapy in Sports: Official Journal of the Association of Chartered Physiotherapist in Sports Medicine*.2014;15(2):82-90.
5. Redmon JM, Gupta A, Stake CE, Hammarstedt JE, et al. Clinical results of hip arthroscopy for labral tears: a comparison between intraoperative platelet-rich plasma and bupivacaine. *Journal Of Arthroscopic & Related Surgery*. 2015;31(3):445-453.
6. InterQual® Level of Care Criteria 2015, Acute Care Adult, McKesson Health Solutions, LLC; 2015 INPT-14.
7. Hufeland M, Kruger D, Haas N, Perka C, Schroder J. Arthroscopic treatment of femoroacetabular impingement shows persistent clinical improvement in the mid-term. *Arch Ortho Trauma Surg*. 2016;136:687-691.
8. Horner N, Ekhtiari S, Simunovic N, Safran M, et al. Hip arthroscopy in patients Age 40 or older: A Systematic Review. *Arthroscopy*. 2016 Sept;1-12.
9. Stephan P, Maarten R, Mathijssen N, Hannink G, Bloem R. Developing a prediction model for the functional outcome after hip arthroscopy. *BMC Musculoskeletal Disorders*. 2018;19:122.
10. Griffin D, Dickenson E, Wall P, et al. Hip arthroscopy versus best conservative care for the treatment of femoroacetabular impingement syndrome (UK FASHIoN): a multicenter randomized trial. *Lancet*. 2018;391:2225-35.