

References

X-27

1. Langhans L, Vejborg TS, Vejborg I, Kroman N. Marking of non-palpable changes in breast tissue. *Ugeskr Laeger*. 2012;174(34):1891-1894.
2. Dahabreh IJ, Wieland LS, Adam GP, Halladay C, Lau J, Trikalinos T.A. Core needle and open surgical biopsy for diagnosis of breast lesions: An update to the 2009 report. Comparative effectiveness review No. 139. (Prepared by the Brown Evidence-based Practice Center under Contract 290-2012-00012-I.) AHRQ Publication No. 14-EHC040-EF. Rockville, MD: Agency for Healthcare Research and Quality. September 2014.
3. Mathenge E, Dean C, Clements D, et al. Core needle biopsy of breast cancer tumors increases distant metastases in a mouse model. *Neoplasia*. 2014;16(11):950-960.
4. Kim E, Hankins A, Cassity J, et al. Isolated radial scar diagnosis by core-needle biopsy: Is surgical excision necessary? *Springer Plus*. 2016;5:398.
5. Siu A. U.S. Preventive Services Task Force. Screening for breast cancer: U.S. Preventive Services Task Force recommendation statement. *Ann of Intern Med*. 2016;164(4):279-296.
6. National Comprehensive Cancer Network (NCCN). Clinical practice guideline: Breast cancer. Version 2.2016.
7. Esen G, Tutar B, Uras C, Calay Z, İnce Ü, Tutar O. Vacuum-assisted stereotactic breast biopsy in the diagnosis and management of suspicious microcalcifications. *Diagn Interv Radiol*. 2016;22(4):326-333.
8. Gümüş H, Mills P, Fish D, et al. Predictive factors for invasive cancer in surgical specimens following an initial diagnosis of ductal carcinoma in situ after stereotactic vacuum-assisted breast biopsy in microcalcification-only lesions. *Diagn and Interv Radiol*. 2016;22(1):29-34.